

Conformal field theory and edge excitations for the principal series of quantum Hall fluids

V. Pasquier and D. Serban

Service de Physique Théorique, CE-Saclay, F-91191 Gif-Sur-Yvette, France.

(February 1, 2008)

Motivated by recent experimental results, we reconsider the theory of the edge excitations for the fractional Hall effect at filling factors $\nu = p/(2np+1)$. We propose to modify the standard $u(1) \otimes su(p)$ edge theory for this series by introducing twist fields which change the boundary conditions of the bosonic fields. This has the effect of removing the conserved charges associated to the neutral modes while keeping the right statistics of the particles. The Green function of the electron in presence of twists decays at long distance with an exponent varying continuously with ν .

The low energy physics of the quantum Hall samples is governed by excitations located at the edge, the bulk excitations being gapped [1]. Recently, it became possible to test the nature of these excitations through experiments involving tunneling conductance measurements [2,3]. Therefore, it is important to have a reliable theoretical description of the edge physics. Wen [4] showed that for filling factors $\nu = 1/(2n+1)$, the edge can be described as a chiral Luttinger liquid. This theory was extended [5] to filling fractions of the form $\nu = p/(2np+1)$, where it involves p independent chiral bosons. The appearance of the quantum Hall effect at these filling fractions is understood in terms of the hierarchical construction [6,7], or more intuitively using the composite fermions picture [8]: the electrons trap an even number $(2n)$ of quanta of flux, see a reduced magnetic field and show their own quantum Hall effect at integer effective filling fraction p . The p effective Landau levels are thought to be at the origin of the p components of the corresponding field theory [9]. It was shown [5] that the edge theory is a conformal field theory with affine symmetry $u(1) \otimes su(p)$. A different version of the conformal field theory on the edge, based on the $W_{1+\infty}$, was proposed [10].

The experiments [2,3] have tested the Luttinger liquid nature of the edge by measuring the tunneling conductance between an electron gas and a fractional quantum Hall fluid, and found a power law dependence on the voltage, $dI/dV \sim V^{\alpha-1}$. Surprisingly, the exponent α varies linearly with $1/\nu$, instead of being discontinuous, as predicted by the theory [11].

Several attempts were made to conciliate the predictions of the edge theory with the experiments [12–15]. The basic problem to be solved concerns the electron's Green function. The electron is a fermion, so the Green function has to be odd under the exchange of the fermion coordinates. On the other hand, to explain the observed dependence of the conductance, the Green function should behave as $t^{-1/\nu}$ at large time separations. The two requirements seem to be contradictory, at least in the context of conformal field theory for the edge. The

second requirement is fulfilled if only the mode carrying the electric charge propagates. Different mechanisms were proposed to suppress the contribution of the neutral modes, otherwise necessary to reestablish the fermionic electron statistics.

In this paper, we show that a consistent mechanism for the suppression of the neutral modes exists in the framework of conformal field theory. We consider a modification the standard p -boson conformal field theory [5] by introducing twist operators. They create branch cuts for the chiral boson fields, change their boundary conditions, and therefore interpolate between sectors of the Hilbert space with different quasi-periodicity properties. Twist operators were considered in the context of conformal field theory on Riemann surfaces [17,18] and string theory on orbifolds [19]. The first effect of twists is to break the global $su(p)$ symmetry, therefore removing the charges associated to the neutral sector, the $su(p)$ spin index being replaced by a branch index. Second, the presence of twist insertions in the electron Green function modify their behavior, depending on the twist coordinates. If a twist is introduced (removed) at $t = -\infty$ ($t = \infty$), then, in the thermodynamic limit we retrieve the same behavior of the Green function as in ref. [11]. In the limiting case where twists are glued to the electrons, the neutral modes are completely suppressed and the Green function behaves as $t^{-1/\nu}$ at time separation t .

To trace back the significance of the twists, in the last section we investigate the properties of the bulk wave functions of free electrons in magnetic field at filling factor $\nu = p/(2np+1)$. Since at this filling factor there are $2n+1/p$ flux quanta per particle, the quantization of the system in a finite geometry (on the torus, for example) can be done consistently only by considering p -valued wave functions. The Hilbert space splits into p quasi-periodicity sectors, which will survive even after introducing an interaction potential. Reducing the electron density to a one-dimensional density on the edge, we retrieve a bosonic fields with twisted boundary conditions. We are lead to conclude that the presence of the twists

in the edge field theory reflect the particular structure of the Hilbert space at the filling factors $\nu = p/(2np + 1)$.

Conformal Field Theory for twisted bosons. Let us first briefly present the basic ingredients of the standard theory of the edge excitations, from the point of view of a conformal field theory [16]. We use the complex coordinate $z = e^{\epsilon - ix}$, in which the surfaces of equal times are circles (radial quantization). Since the excitations at one edge are chiral, we consider only the holomorphic sector of the theory.

Define p bosonic fields by :

$$\varphi_k(z) = \varphi_0^{(k)} - ia_0^{(k)} \ln z + i \sum_{m \neq 0, m \in \mathbf{Z}} \frac{a_m^{(k)} z^{-m}}{m}, \quad \text{with} \\ [a_m^{(k)}, a_{m'}^{(l)}] = m \delta_{m+m', 0} \delta_{k,l}, \quad [\varphi_0^{(k)}, a_0^{(l)}] = i \delta_{k,l}. \quad (1)$$

The bosonic modes $a_m^{(k)}$ with $m < 0$ play the role of creation operators. The fundamental object insuring the conformal invariance is the stress-energy tensor:

$$T(z) = -\frac{1}{2} \sum_{l=0}^{p-1} : (\partial \varphi_l(z))^2 :$$

with conformal anomaly, or central charge $c = p$. The double dots represent normal ordering with respect to the bosonic operators. There are p conserved charges $a_0^{(k)}$, corresponding to the conserved currents $\partial \varphi_k(z)$.

The charged excitations in the spectrum, electrons and quasi-particles, are created by exponentials of bosonic fields (vertex operators). Electrons and quasi-particles are characterized by the dual parameters β and β' , defined by:

$$\beta = 2n, \quad p\beta' + 1 = 1/(p\beta + 1).$$

For the standard theory of edge excitations [9], the electron and quasi-particle operators are:

$$\Psi_e^{(k)}(z) =: e^{i \sum_j v_j^k \varphi_j(z)} :, \quad \Psi_q^{(k)}(z) =: e^{i \sum_j s_j^k \varphi_j(z)} :$$

with $\sum_{j=0}^{p-1} v_j^k v_j^l = \beta + \delta_{kl} \equiv K_{kl}$, $\sum_{j=0}^{p-1} s_j^k v_j^l = \delta_{kl}$ and $\sum_{j=0}^{p-1} s_j^k s_j^l = \beta' + \delta_{kl} = (K^{-1})_{kl}$. The vectors v^k generate the p dimensional charge lattice of a $u(1) \otimes su(p)$ algebra, characterized by the $p \times p$ matrix K [7,9] and the vectors s^k generate the dual lattice. The index (k) of the vertex operators is a $su(p)$ spin index. Properties as the statistics or the dimension of the operators can be deduced from the product of two operators :

$$\Psi_e^{(k)}(z) \Psi_e^{(l)}(w) = (z-w)^{\beta + \delta_{kl}} : \Psi_e^{(k)}(z) \Psi_e^{(l)}(w) :.$$

The electron Green function can also be determined :

$$\langle \Psi_e^{(k)\dagger}(z) \Psi_e^{(k)}(w) \rangle = (z-w)^{-\beta-1},$$

leading to the exponent for the conductance $\alpha = \beta + 1$. Quasi-particle have similar properties, with β replaced by β' .

Let us now allow for the presence of twists, changing the boundary conditions of the bosonic fields. Twists appeared in the conformal field theory in connection with the Ising model [18], where they represent the spin operators, and in string theory [17,19], where they were used to introduce branch point singularities in the background. Here, we use $p-1$ twist operators $\sigma_1, \dots, \sigma_{p-1}$ which introduce branch points of order p for the bosonic fields [19]. We make the change of variable:

$$\tilde{\varphi}_l(z) = \frac{1}{\sqrt{p}} \sum_k \omega_p^{lk} \varphi_k(z),$$

with $\omega_p = e^{2\pi i/p}$. In terms of these fields, the monodromy conditions around the singularities introduced by the twists are given by :

$$\partial \tilde{\varphi}_l(z) \sigma_k(0) = z^{-\alpha/p} \tau_{l,k}(0) + \dots,$$

where $\alpha = lk \bmod p$, the dots indicate less singular terms and the fields $\tau_{l,k}(z)$ are called excited twist fields. The twists themselves obey the following algebra:

$$\sigma_k(z) \sigma_l(0) = (z-w)^\delta \sigma_r(0) + \dots,$$

with $r = k + l \bmod p$, $\sigma_0 \equiv 1$ and δ is a number which can be determined by dimensional analysis. A twist operator interpolates between two different sectors of the Hilbert space.

First, we will consider the simpler case when there is one twist σ_1 located at $z = 0$ and one σ_{p-1} located at $z = \infty$ (the global twist should be equal to zero). The vacuum in this twisted sector is $|\sigma_1\rangle = \sigma_1(0)|0\rangle$, on which the p bosonic fields have the following mode decomposition :

$$\partial \tilde{\varphi}_l(z) = -i \sum_{m \in \mathbf{Z} + l/p} a_m z^{-m-1}, \\ \text{with} \quad [a_m, a_{m'}] = m \delta_{m+m', 0}, \quad (2)$$

Here, the bosonic modes a_m are indexed by a fractional number $m \in \mathbf{Z}/p$, the ones with $m < 0$ representing the creation operators. Note that the commutation relations naturally couple the field $\tilde{\varphi}_l(z)$ to $\tilde{\varphi}_{p-l}(z)$.

The holomorphic component of the stress energy tensor in the sector twisted by σ_1 is:

$$\tilde{T}(z) = -\frac{1}{2} : ((\partial \tilde{\varphi}_0(z))^2 + \partial \tilde{\varphi}_1(z) \partial \tilde{\varphi}_{p-1}(z) + \dots + \\ + \partial \tilde{\varphi}_{p-1}(z) \partial \tilde{\varphi}_1(z)) : + \frac{p^2 - 1}{24pz^2}. \quad (3)$$

Note that the stress energy tensor is single valued around $z = 0$, although the constituting bosons are not. The last term in $T(z)$ gives the dimension of the twist operator σ_1 , $\Delta_1 = (p^2 - 1)/24p$. Similarly, one obtains for the twisted sectors generated by σ_k , $\Delta_k = (p^2 - q^2)/24p$, where q is the greatest common divisor of p and k [19].

From the mode decomposition (2), we deduce that there is no conserved charge associated to the twisted bosons $\tilde{\varphi}_1, \dots, \tilde{\varphi}_{p-1}$. Even if there are conserved currents associated to the twisted sector, they *do not* manifest at the level of global symmetries. The only conserved charge is the electric charge $J_0 = \sqrt{\nu}a_0$, normalized such that the electron charge is 1.

Electron and quasi-particle operators in presence of twist. In the presence of the twist $\sigma_1(0)$ the bosonized form of the electron and the quasi-particle operators is the following:

$$\Psi_{e,q}^{(k)}(z) \sigma_1(0) = z^{-\gamma} \tilde{\Psi}_{e,q}^{(k)}(z) \sigma_1(0), \quad (4)$$

with:

$$\begin{aligned} \tilde{\Psi}_e^{(k)}(z) &= : e^{\frac{i}{\sqrt{p}}(\sqrt{p\beta+1}\tilde{\varphi}_0(z) + \sum_{j=1}^{p-1} \omega_p^{kj} \tilde{\varphi}_j(z))} : , \\ \tilde{\Psi}_q^{(k)}(z) &= : e^{\frac{i}{\sqrt{p}}(\frac{1}{\sqrt{p\beta+1}}\tilde{\varphi}_0(z) + \sum_{j=1}^{p-1} \omega_p^{kj} \tilde{\varphi}_j(z))} : . \end{aligned} \quad (5)$$

The power γ can be computed by dimensional analysis, in the following way. Let $\Delta_e = (\beta + 1)/2$ and $\Delta_q = (1 - \beta/(p\beta + 1))/2$ be the dimensions of the operators $\Psi_{e,q}^{(k)}(z)$ respectively. In the presence of the twist σ_1 , the vertex operators $\tilde{\Psi}_{e,q}^{(k)}(z)$ have the dimensions $\tilde{\Delta}_e = 1/2\nu$ and $\tilde{\Delta}_q = \nu/2p^2$. The reason for this change in the dimensions of the vertex operators is the absence of the zero modes of the fields $\tilde{\varphi}_1, \dots, \tilde{\varphi}_{p-1}$. We obtain $\gamma = \Delta_e - \tilde{\Delta}_e = \Delta_q - \tilde{\Delta}_q = (1 - 1/p)/2$.

As the bosons (2), the vertex operators have a non-trivial monodromy around the point $z = 0$:

$$z \rightarrow ze^{2i\pi} \Rightarrow \Psi_e^{(k)}(z) \rightarrow c_k \Psi_e^{(k-1)}(z).$$

where c_k is a phase factor, and similarly for $\Psi_q^{(k)}(z)$. The index (k) has to be understood modulo p , and it represents now a branch index rather than a spin index. Since the p operators $\Psi_e^{(k)}(z)$ represent different determinations of the same operator, we will drop the branch index in the following.

Next, we are going to compute the Green function of the electron operator, in the presence of two twists at arbitrary positions z_1, z_2 :

$$G(z, w; z_1, z_2) = \frac{\langle \sigma_{p-1}(z_1) \Psi_e^\dagger(z) \Psi_e(w) \sigma_1(z_2) \rangle}{\langle \sigma_{p-1}(z_1) \sigma_1(z_2) \rangle}$$

The short distance ($z \rightarrow w$) singularity of $G(z, w; z_1, z_2)$ is given by $(z-w)^{-\beta-1}$. From the behavior of the correlation function under conformal transformations, we have:

$$G(z, w; z_1, z_2) = (z-w)^{-\beta-1} F(x), \quad x = \frac{(z-z_1)(w-z_2)}{(z-z_2)(w-z_1)}$$

with $F(x)$ some function of the cross ratio x , which can be determined in the limit $z_1 \rightarrow \infty, z_2 \rightarrow 0$ by using the bosonization (5):

$$F(x) = x^{-\frac{1}{2}(1-\frac{1}{p})} \frac{x-1}{x^{1/p}-1}.$$

Since $F(x) = F(1/x)$, the Green function $G(z, w; z_1, z_2)$ is odd under exchange of z and w . It is possible to remove the short distance singularity in the product of electron and twist operator, allowing to bring them eventually to the same point:

$$: \Psi_e(z) \sigma_1(w) : \equiv (z-w)^\gamma \Psi_e(z) \sigma_1(w).$$

Bringing the electron and the twist together cancels the neutral modes, as can be seen from the corresponding Green function:

$$\begin{aligned} G'(z, w; z, w) &= \frac{\langle : \sigma_{p-1}(z) \Psi_e^\dagger(z) : : \Psi_e(w) \sigma_1(w) : \rangle}{\langle \sigma_{p-1}(z) \sigma_1(w) \rangle} \\ &= (z-w)^{-\beta-1/p} = (z-w)^{-1/\nu}. \end{aligned}$$

We infer that this Green function correspond to the insertion of charge only into the system. If the same mechanism is applied to the quasi-particle excitations, the exponent for the quasi-particle Green function will be given by $Q^2/\nu = \nu/p^2$.

For comparison, if the twists are located at $z_1 = \infty, z_2 = 0$ (corresponding to $t_1 = \infty, t_2 = -\infty$), the Green function is:

$$\begin{aligned} G'(z, w; \infty, 0) &= \frac{\langle : \sigma_{p-1}(\infty) \Psi_e^\dagger(z) : : \Psi_e(w) \sigma_1(0) : \rangle}{\langle \sigma_{p-1}(\infty) \sigma_1(0) \rangle} \\ &= (z-w)^{-\beta} (z^{1/p} - w^{1/p})^{-1}. \end{aligned}$$

In the thermodynamic limit $t \ll L/v \equiv 2\pi$, where t is the time separation between the points z and w , this last Green function behaves like $t^{-\beta-1}$ and the effect of the twist cannot be detected.

Electrons on the torus at $\nu = p/(p\beta + 1)$. In order to understand better the origin of the twists in the edge theory we go back and study the problem of N electrons at filling factor $\nu = p/(p\beta + 1)$. For the sake of simplicity, we use a torus of periods L_x and L_y and area $S = L_x L_y$,

In the absence of Coulomb interactions, the Hamiltonian is a sum of one-particle terms of the type:

$$H_1 = \frac{1}{2m_e} (\vec{p} - e\vec{A})^2$$

The magnetic translations $T(\vec{R}) = e^{\frac{i}{\hbar} \vec{R}(\vec{p} + e\vec{A})}$ commute with the Hamiltonian, but do not mutually commute. The magnetic translations along the two periods of the torus, T_x and T_y obey the commutation relation:

$$T_x T_y = e^{2\pi i \frac{BS}{\phi_0}} T_y T_x,$$

$\phi_0 = h/e$ being the flux quantum. The number of flux quanta through the system $BS/\phi_0 = N/\nu = N(\beta + 1/p)$ is in general fractional, so that the two translations cannot be diagonalized simultaneously and one cannot find a basis of periodic one-particle wave functions. Put differently, the number of states per Landau level, $N/\nu =$

$N(\beta + 1/p)$, should be fractional. Since $T_x^p T_y = T_y T_x^p$, the wave functions could be rendered periodic on a p -cover of the torus, with periods pL_x and L_y for example. To preserve the filling factor, we have to consider all the wave functions periodic on the p -cover of the initial torus and fill them with pN electrons.

Let us construct explicitly the one-particle wave functions on the torus from the wave functions on the plane, in the gauge $A_x = -By$, $A_y = 0$. For $b = eB/\hbar > 0$, the wave functions in the lowest Landau level are:

$$\phi_m(x, y) = C e^{-b(y-2\pi m/bL_x)^2/2} e^{-2\pi i m x/L_x},$$

with C a normalization constant. To render these functions periodic over L_y , one takes:

$$\Phi_m(x, y) = \sum_{M \in \mathbf{Z}} \phi_{m+MN(\beta+1/p)}(x, y).$$

If $m \in \mathbf{Z}/p$, the resulting function will be periodic over pL_x . Their behavior under translations by L_x depend on the value of N . We identify p sectors in the Hilbert space, indexed by $k = 0, \dots, p-1$, where the number of electrons is $N = k \bmod p$. For simplicity, we consider in the following the sector $k = 1$. Then, the wave functions $\Phi_m(x, y)$ decompose into p sectors:

$$\begin{aligned} \Phi_m(x, y) &= \sum_{j=0}^{p-1} \Phi_m^{(j)}(x, y), \\ \Phi_m^{(j)}(x + L_x, y) &= \omega_p^j \Phi_m^{(j)}(x, y). \end{aligned}$$

With $m = 1/p, 2/p, \dots, N/\nu$, they constitute a $N(p\beta + 1)$ dimensional basis of the lowest Landau level on the torus of area $pL_x L_y$. A small perturbation potential, periodic on the initial torus, will preserve the quasi-periodicity sectors, since the matrix elements of the potential between two different periodicity sectors vanish.

To deduce the implications of the quasi-periodicity for the edge excitations, we consider the situation where two (sharp) edges are created at $y = 0$ and $y = L$, by introducing some confining potential. For simplicity, we take $L, L_y \rightarrow \infty$ so that the wave functions are equal to $\phi_m(x, y)$, normalized by $C = (pL_x \sqrt{\pi/b})^{-1/2}$. The electron operator reads:

$$\Psi(x, y) = \sum_{m \in \mathbf{Z}/p} \phi_m(x, y) c_m.$$

Following the procedure used in ref. [13], we define a one-dimensional density operator, by integrating the two dimensional density operator from $y = -\infty$ to some cutoff $y = Y$ in the bulk:

$$\rho(x) = \int_{-\infty}^Y \Psi^\dagger(x, y) \Psi(x, y) dy.$$

We obtain a one-dimensional density operator which is not periodic in x :

$$\begin{aligned} \rho(x) &= \sum_{l=0}^{p-1} \rho^{(l)}(x), \quad \rho^{(l)}(x + L_x) = \omega_p^l \rho^{(l)}(x), \\ \rho^{(l)}(x) &= \sum_{m \in \mathbf{Z} + l/p} e^{2\pi i m x/L_x} \rho_m. \end{aligned}$$

Replacing the fermion number with its average in the ground state, $\langle 0 | c_m^\dagger c_{m'} | 0 \rangle = \nu \delta_{m, m'}$, one can show that the low modes of the one-dimensional density operator, $|m| \ll L_x \sqrt{b}/2\pi$, obey bosonic commutation relations:

$$[\rho_m, \rho_{m'}] = \nu m \delta_{m+m', 0}, \quad m, m' \in \mathbf{Z}/p.$$

With ρ_m replaced by $\sqrt{\nu} a_m$, these are the commutation relations of the bosonic fields in the sector twisted by $\sigma_1(2)$. By this identification, $\rho_0 = \sqrt{\nu} a_0$ represents the charge operator. Compared to Zülicke and MacDonald [13], we obtain non-integer modes for the density operator. They correspond to the neutral modes which restore the fermionic statistics of the electron operator.

In conclusion, we have shown that the requirement that only the charge mode contributes to the tunneling conductance can be formulated in a consistent way in the language of conformal field theory. Our conclusions essentially agree with that of references [13–15], the difference coming from the treatment of the neutral modes. We argue that at filling fraction $\nu = p/(2np + 1)$, there are $2n + 1/p$ flux quanta per electron, and the structure of the Hilbert space for the bulk electrons induced by the nonintegrality of the flux is translated on the edge by the help of the twist operators. The twists are able to account for the absence of the neutral modes contribution from the correlation functions, being an alternative to setting the neutral mode velocity to zero from the beginning [14]. More insight about the nature of the quasi-particle excitations come from an analysis of the bulk. It is possible to obtain, in a Landau-Ginsburg-Chern-Simons approach for the bulk, vortex-like solutions which carry only one charge and show a multivalued structure reminiscent from twists [23].

As a technical remark, we point out that the relation between the conformal field theory with $u(1) \otimes su(p)$ symmetry and its twisted version borrows a lot of similarity to the relation between two integrable models, namely the Calogero-Sutherland model with $su(p)$ symmetry and a certain limit of the Ruijsenaars-Schneider model, relation uncovered by Uglov [20]. The bosonization of the charged excitations in the twisted sectors is suggested by this correspondence (see also [21]).

We would like to thank M. Bershadski, A. Cappelli, E. Fradkin, C. Glatli, K. Imura and N. Sandler for useful discussions. After completing this work, we became aware of ref. [22] where twist operators were also used to determine the ground state wave function.

- [1] B. Halperin, Phys. Rev. B **25**, 2185 (1982);
- [2] A.M. Chang, L.N. Pfeiffer and K.W. West, Phys. Rev. Lett. **77**, 2538 (1996);
- [3] M. Grayson, D.C. Tsui, L.N. Pfeiffer, K.W. West and A.M. Chang, Phys. Rev. Lett. **80**, 1062, (1998);
- [4] X.G. Wen, Phys. Rev. B **41**, 12838 (1990);
- [5] J. Fröhlich and A. Zee, Nucl. Phys. B **364**, 517 (1991);
- [6] F.D.M. Haldane, Phys. Rev. Lett. **51**, 605 (1984); B. Halperin, Phys. Rev. Lett. **52**, 1583 (1984);
- [7] N. Read, Phys. Rev. Lett. **65**, 1502 (1990);
- [8] J.K. Jain, Phys. Rev. Lett. **63**, 199, (1989); Phys. Rev. B **41**, 7653, (1990).
- [9] X.G. Wen and A. Zee, Phys. Rev. B **44**, 274 (1991);
- [10] A. Cappelli, C.A. Trugenberger and G. Zemba, Nucl. Phys. B **448**, 470, (1995).
- [11] C.N. Kane and M.P.A. Fisher, Phys. Rev. B **51**, 13449 (1995);
- [12] A.V. Shytov, L.S. Levitov and B.I. Halperin, Phys. Rev. Lett. **80**, 141 (1998);
- [13] U. Zülicke and A.H. MacDonald, Phys. Rev. B **60**, 1837 (1999); S. Conti and G. Vignale, J. Phys. C **50**, 779 (1998);
- [14] D.H. Lee and X.G. Wen, cond-mat/9809160 (unpublished);
- [15] E. Fradkin and A. Lopez, Phys. Rev. B **59**, 15323 (1999);
- [16] P. Di Francesco, P. Mathieu and D. Sénéchal, Conformal Field Theory, Springer, 1997;
- [17] V. Knizhnik, Comm. Math. Phys. **112**, 567 (1987);
- [18] A.I.B. Zamolodchikov, Nucl. Phys. B **285**, 481, (1987);
- [19] M. Bershadsky and A. Radul, Int. J. Mod. Phys. A **2**, 165, (1987); L. Dixon, D. Friedan, E. Martinec and S. Shenker, Nucl. Phys. B **282**, 13, (1987);
- [20] D. Uglov, Comm. Math. Phys. **191**, 663 (1998);
- [21] M.C. Bergère, J. Math. Phys. **41**, 7234, (2000);
- [22] G. Cristofano, G. Maiella, V. Marotta, Mod. Phys. Lett. A **15**, 1679, (2000);
- [23] D. Serban, Phys. Lett. A (to be published).